



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

which is the required formula.¹

(b) Since a and d are positive acute angles, the last term cannot become negative and is zero only when $x = 0$. Therefore d will have a minimum value when $x = 0$, so that $r = r' = \frac{1}{2}a$, and $i = i'$, and the complete ray is symmetrically situated with respect to the prism.

(c) For this minimum value of d , say d_0 , we have the classical laboratory formula

$$n = \frac{\sin \frac{1}{2}(a + d_0)}{\sin \frac{1}{2}a}$$

2879 [1921, 89]. Proposed by E. J. OGLESBY, Washington Square College.

Given the values of $U_{5;9}$, $U_{5;10}$, $U_{5;11}$, $U_{6;9}$, $U_{6;10}$, $U_{6;11}$, $U_{7;9}$, $U_{7;10}$, $U_{7;11}$ where $U_{h;k} = \sqrt{hk}$, find the value of $U_{6.2;9.3}$ by interpolation.

SOLUTION BY J. B. REYNOLDS, Lehigh University.

From the first three given terms we find by the method of differences² $U_{5;9.3} = 6.8190$, from the next three $U_{6;9.3} = 7.4698$ and from the last three $U_{7;9.3} = 8.0685$. Using these three values then as a series by the same method we find $U_{6.2;9.3} = 7.5937$.

Also solved by H. N. CARLETON.

2881 [1921, 89]. Proposed by E. B. ESCOTT, Oak Park, Ill.

If, in the polynomial $X^3 - 2$, we substitute $x^2 + x - 4$ for X , the given expression can be factored, that is, $X^3 - 2 \equiv (x^3 + 3x^2 - 3x - 11)(x^3 - 6x + 6)$. Find a substitution for X so that the polynomial $X^3 + pX^2 + qX + r$ may be factored.

SOLUTION BY THE PROPOSER.

Let

$$X^3 + pX^2 + qX + r = (X - b)(X + c)^2 - a^2(X - d)^2. \quad (1)$$

Expanding the second member and equating coefficients of like powers of X , we get

$$a^2 + b - 2c + p = 0, \quad 2a^2d - 2bc + c^2 - q = 0, \quad \text{and} \quad a^2d^2 + bc^2 + r = 0.$$

Eliminating b and d and solving for a^2 we get

$$a^2 = \frac{(3c^2 - 2pc + q)^2}{4(c^3 - pc^2 + qc - r)}.$$

Therefore, it is necessary and sufficient that³

$$c^3 - pc^2 + qc - r = n^2,$$

and we get by substitution

$$a = \frac{3c^2 - 2pc + q}{2n}, \quad d = -\frac{c^3 - qc + 2r}{3c^2 - 2pc + q}, \quad \text{and} \quad b = -a^2 + 2c - p.$$

¹ We might say

$$\begin{aligned} \frac{1}{n^2} &= \frac{\sin^2 \frac{1}{2}a \cos^2 x}{\sin^2 \frac{1}{2}(a + d)} + \frac{\cos^2 \frac{1}{2}a \sin^2 x}{\cos^2 \frac{1}{2}(a + d)} \\ &= \frac{\sin^2 \frac{1}{2}a}{\sin^2 \frac{1}{2}(a + d)} + \left[\frac{\cos^2 \frac{1}{2}a}{\cos^2 \frac{1}{2}(a + d)} - \frac{\sin^2 \frac{1}{2}a}{\sin^2 \frac{1}{2}(a + d)} \right] \sin^2 x \\ &= \frac{\sin^2 \frac{1}{2}a}{\sin^2 \frac{1}{2}(a + d)} + \frac{4 \sin \frac{1}{2}d \sin(a + \frac{1}{2}d) \sin^2 x}{\sin^2(a + d)} \end{aligned}$$

and use this formula for (b) and (c) instead of the formula in the text, d being a minimum when $[\sin^2 \frac{1}{2}a]/[\sin^2 \frac{1}{2}(a + d)]$ is a maximum.—EDITORS.

² See 1921, 330.

³ Thus it seems to be a necessary part of the hypothesis that there is a rational number c that will make the given polynomial equal to minus the square of a rational number.

Letting f denote the polynomial, if $f(-c) = -n^2$ we can write

$$f(X) = (X + c)^2 \left[X - 2c + p + \left(\frac{f'(-c)}{2n} \right)^2 \right] - \left[\frac{f'(-c)}{2n} (X + c) - n \right]^2,$$

which we can make the difference of two squares by putting $X - 2c + p + \left(\frac{f'(-c)}{2n} \right)^2 = (x + t)^2$.

—EDITORS.

Thus, a , b , c , and d are known. The right-hand member of (1) will be the difference of two squares, and may be factored if we put $X - b = (x + t)^2$; i.e., $X = (x + t)^2 + b$.

In the example given, $p = q = 0$, $r = -2$. The value $c = -1$ gives $n = 1$; whence, $a = 3/2$, $d = 5/3$, $b = -17/4$. $X^3 - 2 = (X + \frac{1}{4})(X - 1)^2 - \frac{1}{4}(3X - 5)^2$. Letting $X + \frac{1}{4} = (x + \frac{1}{2})^2$; i.e., $X = x^2 + x - 4$, and substituting, we have $X^3 - 2 = [(x + \frac{1}{2})(x^2 + x - 5) + \frac{1}{2}(3x^2 + 3x - 17)][(x + \frac{1}{2})(x^2 + x - 5) - \frac{1}{2}(3x^2 + 3x - 17)] = (x^3 + 3x^2 - 3x - 11)(x^3 - 6x + 6)$.

2885 [1921, 139].

If A, B, C, X, Y are given collinear points, construct Z so that $\{ABCX\} + \{ABCY\} = \{ABCZ\}$, where $\{ABCX\}$ denotes the cross-ratio of the points A, B, C, X . [From the *Math. Tripos Exam.*, Cambridge, Eng., 1905.]

I. SOLUTION BY GERTRUDE I. MCCAIN, Westminster College, New Wilmington, Pa.

If $\{ABCX\} + \{ABCY\} = \{ABCZ\}$, then $\frac{AB}{BC} : \frac{AX}{XC} + \frac{AB}{BC} : \frac{AY}{YC} = \frac{AB}{BC} : \frac{AZ}{ZC}$. Dividing by $\frac{AB}{BC}$, $\frac{XC}{AX} + \frac{YC}{AY} = \frac{ZC}{AZ}$. But $XC = -CX = -(AX - AC)$, $YC = -CY = -(AY - AC)$, and $ZC = -CZ = -(AZ - AC)$. Making the substitutions and adding 2 to each side of the equation, we have $\frac{AC}{AX} + \frac{AC}{AY} = \frac{AC}{AZ} + 1$, or $\frac{1}{AX} + \frac{1}{AY} = \frac{1}{AZ} + \frac{1}{AC}$. Let $AX = \cot \theta$, $AY = \cot \varphi$, $AC = \cot \psi$, and $AZ = \cot \tau$. Considering AX , AY and AC as line functions, construct the angles θ , φ and ψ , using a circle with radius unity. Also construct the tangents of θ , φ , and ψ as line functions using the same circle.

It is now possible to lay off $\tan \theta$ and $\tan \varphi$ as lengths on a line and from their sum subtract $\tan \psi$.

But $\frac{1}{AX} + \frac{1}{AY} - \frac{1}{AC} = \frac{1}{AZ} = \tan \tau$. $\cot \tau$ or AZ may then be constructed as a line function, and the point Z located on the line with A, B, C, X, Y .

II. SOLUTION BY OTTO DUNKEL, Washington University.

After showing that $1/AX + 1/AY = 1/AZ + 1/AC$, it should be observed that this result shows that the two sets of points A, X, Y and A, Z, C have a common fourth harmonic point K . Hence K can be constructed from the first set and then Z as the harmonic conjugate of C with respect to A and K , by the quadrilateral construction, thus giving a non-metrical construction. This would then give a projective definition of the sum of two cross-ratios. This result may be obtained by use of abridged notation as follows:

If $\alpha = 0$, $\beta = 0$, $\gamma = 0$, $\xi = 0$, $\eta = 0$ are the equations of the points A, B, C, X, Y , then the expressions to the left in these equations may be chosen so that

$$\beta = b\alpha + \gamma, \quad \xi = x\alpha + \gamma, \quad \eta = y\alpha + \gamma,$$

and then $\{ABCX\} = x/b$, $\{ABCY\} = y/b$. It follows that $(x - y)\alpha = \xi - \eta$, and if K is the harmonic conjugate of A with respect to X, Y , its equation may be written $(x - y)\kappa = \xi + \eta = 0$ or $(x - y)\kappa = (x + y)\alpha + 2\gamma$. Since $2\gamma = (x - y)\kappa - (x + y)\alpha$, Z , the harmonic conjugate of C with respect to K, A , may have its equation written $2\xi = (x - y)\kappa + (x + y)\alpha = 0$ or $\xi = (x + y)\alpha + \gamma$. Hence $\{ABCZ\} = (x + y)/b = \{ABCX\} + \{ABCY\}$.

Also solved by NATHAN ALTSHILLER-COURT, J. W. CLAWSON, ARTHUR PELLETIER, and F. L. WILMER.

2887 [1921, 139]. Proposed by the late L. G. WELD.

A carpenter's square moves with its outer edges in contact with two round pegs of given equal diameters. Define the locus of the "heel" of the square.

SOLUTION BY J. B. REYNOLDS, Lehigh University.

Let the radius of either peg be r and the distance between their centers $\sqrt{2}a$. Let the axes be taken so that the centers of the circles are at the points $(a, 0)$ and $(0, a)$, and assume that the